



MOLYBDENUM 10%

GUARANTEED ANALYSIS

Molybdenum (Mo) 10%
10% Water Soluble Molybdenum (Mo)

Derived from Sodium Molybdate

NOTICE

The application of fertilizing material containing molybdenum (Mo) may result in forage crops containing levels of molybdenum (Mo) which are toxic to ruminant animals.

PRODUCT DESCRIPTION

Monty's Molybdenum 10% is to be used to prevent and correct molybdenum deficiencies.

DIRECTIONS FOR USE

Foliar Application: Actively growing plants with young leaves absorb foliar applied Monty's Molybdenum 10% most efficiently. Immediate post-irrigation or post-rainfall application will enhance foliar uptake. Thorough coverage and wetting are needed for optimum foliar spray results. Applications should be separated by a minimum of 7-14 days. AVOID foliar applications of this product when plants are under moisture stress. AVOID foliar applications to deciduous tree crops during bloom. Application Rate: 2-5 fluid ounces/acre.

Soil Application: Monty's Molybdenum 10% can be soil applied but should be incorporated. For best results when soil applied, split applications work best. Monty's Molybdenum can be applied alone or in combination with liquid fertilizers. Application Rate: 2-5 fluid ounces/acre; Light Deficiency 10-20 fluid ounces; Moderate Deficiency 20-30 fluid ounces; Heavy Deficiency 30-40 fluid ounces. Fertilization: Apply the recommended amount of product in split applications 1-2 weeks apart throughout the growing season.

Net Contents - 2.5 US Gal (9.46 Liters)

Net Weight - 25 Lbs (11.3 Kg)

10.01 Lbs/Gal at 68° F



8727

F001360

6 36329 00726 2



www.montysplantfood.com
Monty's Plant Food Company
4800 Strawberry Lane
Louisville, KY 40209
800-978-6342

Monty's products are recommended as supplements to your current farming practices and are not recommended to replace the nutrients typically added to the soil. We suggest taking a soil and/or tissue test, have a qualified independent lab analyze the samples to determine nutrient levels in your soils and plants, and then follow the recommendations of a qualified CCA, agronomist or dealer.

When tank mixing any Monty's products with other farm chemicals, always read and follow the label directions of the chemical(s) you are considering and administer a jar test and/or a strip test before adding Monty's to your tank mix. When mixing with other farm chemicals, tank mix Monty's only after diluting the chemical according to label directions. Knowing the chemical's "mode of action" (block photosynthesis) or metal content (copper based) will help determine the compatibility of Monty's.

Storage and Handling

Use within 24 hours of mixing with water, chemicals or other products. Keep container closed when not in use. Avoid freezing. Products containing organic materials may settle and should be shaken or stirred if stored. Products containing living organisms should not be stored in direct sunlight or allowed to freeze; should not be carried over from one year to the next; and should be used during the growing season in which it was purchased. Store in a cool, dry place. Refer to the SDS for other safety and handling information.

Limited Warranty

Monty's warrants that this product meets its manufacturing specifications. If it does not, Monty's will, at its option, replace the product or refund the purchase price. In no event shall Monty's be liable for special, incidental or consequential damages or for damages in the nature of penalties. Monty's shall not be liable in any way for claims resulting from any use of this product which is not in strict accordance with all directions, cautions and warnings on the label or labels. For products containing living organisms, this warranty is limited to the "use by date" on the label.

FOR COMMERCIAL USE ONLY. KEEP OUT OF REACH OF CHILDREN. HARMFUL IF SWALLOWED. AVOID CONTACT WITH SKIN, EYES, OR CLOTHING.

Information regarding the contents and levels of metals in this product is available on the Internet at:
<http://www.aapfco.org/metals.htm>

